

# Docetaxel and cyclophosphamide

## Indication

Adjuvant treatment for breast cancer

## Regimen details

Day	Drug	Dose	Route
1	Cyclophosphamide	600mg/m <sup>2</sup>	IV slow bolus
1	Docetaxel	75mg/m <sup>2</sup>	IV infusion

## Cycle frequency

21 days

## Number of cycles

6 cycles

## Administration

Administer cyclophosphamide first

Cyclophosphamide is administered by slow IV bolus into the arm of a fast-running drip of sodium chloride 0.9%. Cyclophosphamide may also be given as an IV infusion in 250-500mL sodium chloride 0.9% over 30 minutes.

Docetaxel is administered as an IV infusion in 250mL or 500mL (concentration dependent) PVC free sodium chloride 0.9% over 60 minutes.

Patients should be observed closely for hypersensitivity reactions, particularly during the first and second infusions.

Hypersensitivity reactions may occur within a few minutes following the initiation of the infusion of docetaxel and therefore facilities for the treatment of hypotension and bronchospasm must be available.

If hypersensitivity reactions occur, minor symptoms such as flushing or localised cutaneous reactions do not require discontinuation of therapy. The infusion may be temporarily interrupted and when symptoms improve restarted at a slower infusion rate. Severe reactions, such as hypotension, bronchospasm or generalised rash/erythema require immediate discontinuation of docetaxel and appropriate therapy.

Patients who have developed severe hypersensitivity reactions should not be re-challenged with docetaxel.

## Pre-medication

Dexamethasone 8 mg BD (morning and lunchtime) for 3 days starting 24 hours prior to chemotherapy. (Note: Patients must receive 3 doses of dexamethasone prior to treatment).

In the case where 3 doses have not been taken, dexamethasone 16-20mg IV should be administered 30-60 minutes prior to chemotherapy and the remaining 3 oral doses should be taken as normal.

## Emetogenicity

This regimen has mild emetic potential

## Additional supportive medication

Mouthwashes as per local policy

H<sub>2</sub> antagonist or proton-pump inhibitor if required  
 Loperamide if required.  
 Scalp cooling may be offered.  
 Filgrastim or pegfilgrastim should be given, starting on day 2

### Extravasation

Docetaxel is an exfoliant (Group 4)  
 Cyclophosphamide is neutral (Group 1)

### Investigations – pre first cycle

Standard pre-SACT tests

### Investigations - pre subsequent cycles

Investigation	Validity period (or as per local policy)
FBC	96 hours
U+E (including creatinine)	7 days
LFTs	7 days

### Standard limits for administration to go ahead

If blood results not within range, authorisation to administer **must** be given by prescriber/ consultant

Investigation	Limit
Neutrophils	$\geq 1.0 \times 10^9/\text{L}$
Platelets	$\geq 100 \times 10^9/\text{L}$
Creatinine Clearance (CrCl)	$> 20 \text{ mL/min}$
Bilirubin	$\leq 1.0 \text{ ULN}$
AST/ALT	$\leq 1.5 \times \text{ULN}$
Alkaline Phosphatase	$\leq 2.5 \times \text{ULN}$

### Dose modifications

- Haematological toxicity**

If neutrophils  $< 1.0 \times 10^9/\text{L}$  and/or platelets  $< 100 \times 10^9/\text{L}$  delay 1 week or until recovery.

If febrile neutropenia or neutrophils  $< 0.5 \times 10^9/\text{L}$  for more than 1 week consider reduction doses of docetaxel and cyclophosphamide to 80% for future cycles.

- Renal impairment**

CrCl (mL/min)	Cyclophosphamide dose
$> 20$	100%
10-20	75%
$< 10$	50%

There is no data available on the use of docetaxel in severe renal impairment. No modifications required.

- Hepatic impairment**

AST/ALT (x ULN)		Alkaline phosphatase (x ULN)	Docetaxel dose	Cyclophosphamide dose
≤ 1.5	and	< 2.5	100%	100%
> 1.5	or	≥ 2.5- 6	75%	100%
> 3.5	or	≥ 6	Discuss with consultant	Consider dose reduction (discuss with consultant)

If bilirubin > 1.0 x ULN withhold dose (or consultant decision to treat)

- **Other toxicities**

Toxicity	Definition	Docetaxel dose
Peripheral neuropathy	Grade 2	75%
	Grade 3 or 4	Discuss with consultant
Diarrhoea	Grade 3 or 4	1 <sup>st</sup> occurrence – 75%
		2 <sup>nd</sup> occurrence – 60%
Stomatitis	Grade 3 or 4	1 <sup>st</sup> occurrence – 75%
		2 <sup>nd</sup> occurrence – 60%

Any other grade 3 or 4 toxicity- discuss with consultant.

### Adverse effects - for full details consult product literature/ reference texts

- **Serious side effects**

Secondary malignancy  
 Myelosuppression  
 Infusion related reactions  
 Anaphylaxis  
 Teratogenicity  
 Infertility  
 Cardiotoxicity  
 Peripheral neuropathy

- **Frequently occurring side effects**

Diarrhoea  
 Constipation  
 Fatigue  
 Nausea and vomiting  
 Myelosuppression  
 Stomatitis and mucositis  
 Arthralgia and myalgia  
 Alopecia

- **Other side effects**

Fluid retention  
 Deranged liver function  
 Phlebitis  
 Skin toxicity  
 Nail changes  
 Taste disturbances  
 Bladder irritation

### Significant drug interactions – for full details consult product literature/ reference texts

**CYP3A4 Enzyme inducers/inhibitors:** in vitro studies suggest that CYP3A inhibitors (such as ketoconazole, ritonavir, clarithromycin and erythromycin) may raise docetaxel levels, whereas CYP3A inducers (such as rifampicin and barbiturates) may reduce docetaxel levels.

**Warfarin/coumarin anticoagulants:** increased or fluctuating anticoagulant effects. Avoid if possible, consider switching patient to a low molecular weight heparin during treatment or if the patient continues taking an oral anticoagulant monitor the INR at least once a week and adjust dose accordingly.

#### **Cyclophosphamide:**

**Amiodarone:** increased risk of pulmonary fibrosis – avoid if possible

**Clozapine:** increased risk of agranulocytosis – avoid concomitant use

**Digoxin tablets:** reduced absorption – give as liquid form

**Indapamide:** prolonged leucopenia is possible - avoid

**Itraconazole:** may increase adverse effects of cyclophosphamide

**Phenytoin:** reduced absorption - may need to increase dose of phenytoin

**Grapefruit juice:** decreased or delayed activation of cyclophosphamide. Patients should be advised to avoid grapefruit juice for 48 hours before and on day of cyclophosphamide dose.

### Additional comments

Nil

---

### References

Jones S, Holmes FA, O'Shaughnessy J, Blum JL, Vukelja SJ, McIntyre KJ. Et al. Docetaxel with cyclophosphamide is associated with an overall survival benefit compared with doxorubicin and cyclophosphamide: 7 year follow up of US Oncology Research trial 9735. *J Clin Oncol*, 2009; 27:1177-83

---

THIS PROTOCOL HAS BEEN DIRECTED BY DR BEZECNY, CONSULTANT ONCOLOGIST

RESPONSIBILITY FOR THIS PROTOCOL LIES WITH THE HEAD OF SERVICE

DATE May 2025

REVIEW May 2027

VERSION 7

---