

# Policy for Haemorrhoid Surgery

|  |   |
|--|---|
| Ref:   | LSCICB_Clin25   |
| Version:   | 2   |
| Purpose  | This document is part of a suite of policies that the Integrated Care Board (ICB) uses to drive its commissioning of healthcare. Each policy in that suite is a separate public document in its own right but will be applied with reference to other policies in that suite. |
| Supersedes:  | 1.1   |
| Author (inc Job Title):                            | Clinical Policy Group<br>Debbie Lowe - Head of Service – IFR & Clinical Policy  |
| Ratified by:<br>(Name of responsible<br>Committee) | Commissioning Committee   |
| Cross reference to other<br>Policies/Guidance      |   |
| Date Ratified:                                     | 22 April 2026   |
| Date Published and where<br>(Intranet or Website): | April 2026<br>(Website)   |
| Review date:                                       | April 2031  |
| Target audience:                                   | All LSCICB Staff  |

This policy can only be considered valid when viewed via the ICB website or ICB staff intranet. If this document is printed into hard copy or saved to another location, you must check that the version number on your copy matches that of the one published.

| <b>Document control:</b> |                        |   |
|--------------------------|------------------------|---|
| <b>Date:</b>             | <b>Version Number:</b> | <b>Section and Description of Change</b>  |
| March 2019               | V1                     | Policy ratified by Healthier Lancashire and South Cumbria's Joint Committee of Clinical Commissioning Groups (JCCCGs)   |
| July 2022                | V1.1                   | Policy adopted by Lancashire and South Cumbria ICB – references to CCG replaced by ICB where relevant   |
| April 2026               | V2                     | <p>Procedure codes and reference sections updated for accuracy and consistency, with associated updates to the following:</p> <ul style="list-style-type: none"> <li>• Clarified advice provision for all patients and expanded definition of non-operative measures.</li> <li>• Updated wording to reflect Grade 4 haemorrhoids as a distinct classification.</li> <li>• Acknowledged that some patients will proceed directly to surgery where non-surgical options are not appropriate.</li> <li>• Added emphasis on shared decision-making for surgical approach, in line with updated EBI guidance.</li> </ul> |

## **1. Policy**

1.1 All patients with haemorrhoids should be provided with dietary and lifestyle advice.

1.2 The ICB will commission haemorrhoid surgery in the following circumstances:

1.2.1 Persistent grade 1 (rare) or grade 2 haemorrhoids that have not responded to non-operative measures including dietary and lifestyle changes and non-surgical procedures, such as rubber band ligation, injection sclerotherapy, and infrared coagulation, as appropriate

**OR**

1.2.2 Grade 3, grade 4 or combined internal/external haemorrhoids causing persistent pain or bleeding which are recurrent or which are not suitable for non-surgical procedures

**OR**

1.2.3 Large, irreducible external haemorrhoids.

1.3 In cases where there is significant rectal bleeding the patient should be examined internally by a specialist

1.4 When a patient meets the criteria for haemorrhoid surgery, there should be a shared decision-making process to decide what specific surgical procedure is used, taking into account patient choice, haemorrhoid severity, medical comorbidities and local availability.

## **2. Scope and definitions**

2.1 This policy is based on the ICB's Statement of Principles for Commissioning of Healthcare (version in force on the date on which this policy is adopted).

2.2 Haemorrhoid surgery, including haemorrhoidectomy, stapled haemorrhoidopexy and haemorrhoidal artery ligation are surgical procedures to remove haemorrhoids.

2.3 The scope of this policy includes requests for haemorrhoid surgery for patients (both adults and children) who have:

- failed to respond to non-operative measures
- severe haemorrhoids
- recurrent haemorrhoids
- haemorrhoids with a symptomatic external component

2.4 The scope of this policy does not include non-surgical options for the management of haemorrhoids, such as dietary measures, rubber band ligation or sclerotherapy.

2.5 The ICB recognises that a patient may have certain features, such as

- having haemorrhoids
- wishing to have a service provided for their haemorrhoids
- being advised that they are clinically suitable for haemorrhoid surgery; and
- being distressed by their haemorrhoids, and by the fact that they may not meet the criteria specified in this commissioning policy.

Such features place the patient within the group to whom this policy applies and do not make them exceptions to it.

2.6 For the purpose of this policy the ICB defines haemorrhoids as swellings in the rectum and/or anus containing enlarged blood vessels.

2.7 Severe haemorrhoids are defined as:

- recurrent grade 3 or grade 4 combined internal/external haemorrhoids with persistent pain or bleeding; or
- irreducible and large external haemorrhoids

### **3. Appropriate Healthcare**

3.1 The purpose of haemorrhoid surgery is normally to prevent complications of haemorrhoids and alleviate their negative impact on daily life.

3.2 The ICB regards the achievement of this purpose as according with the Principle of Appropriateness. Therefore, this policy does not rely on the principle of appropriateness. Nevertheless, if a patient is considered exceptional in relation to the principles on which the policy does rely, the ICB may consider the principle of appropriateness in the particular circumstances of the patient in question before confirming a decision to provide funding.

### **4. Effective Healthcare**

4.1 This policy relies on the criterion of effectiveness in that the ICB considers that the potential risks associated with the surgical management of asymptomatic or minimally symptomatic haemorrhoids outweigh the potential benefits.

### **5. Cost Effectiveness**

5.1 The ICB does not call into question the cost-effectiveness of haemorrhoid surgery and therefore this policy does not rely on the Principle of Cost-Effectiveness. Nevertheless, if a patient is considered exceptional in relation to the principles on which the policy does rely, the ICB may consider whether the treatment is likely to be Cost Effective in this patient before confirming a decision to provide funding.

### **6. Ethics**

6.1 The ICB does not call into question the ethics of haemorrhoid surgery and therefore this policy does not rely on the Principle of Ethics. Nevertheless, if a patient is considered exceptional in relation to the principles on which the policy does rely, the ICB may consider whether the treatment is likely to raise ethical concerns in this patient before confirming a decision to provide funding.

### **7. Affordability**

7.1 The ICB does not call into question the affordability of haemorrhoid surgery and therefore this policy does not rely on the Principle of Affordability. Nevertheless, if a patient is considered exceptional in relation to the principles on which the policy does rely, the ICB may consider whether the treatment is likely to be affordable in this patient before confirming a decision to provide funding.

## 8. Exceptions

- 8.1 The ICB will consider exceptions to this policy in accordance with the Policy for Considering Applications for Exceptionality to Commissioning Policies.
- 8.2 In the event of inconsistency, this policy will take precedence over any non-mandatory NICE guidance in driving decisions of this ICB. A circumstance in which a patient satisfies NICE guidance but does not satisfy the criteria in this policy does not amount to exceptionality.

## 9. Force

- 9.1 This policy remains in force until it is superseded by a revised policy or by mandatory NICE guidance relating to this intervention, or to alternative treatments for the same condition.
- 9.2 In the event of NICE guidance referenced in this policy being superseded by new NICE guidance, then:
- If the new NICE guidance has mandatory status, then that NICE guidance will supersede this policy with effect from the date on which it becomes mandatory.
  - If the new NICE guidance does not have mandatory status, then the ICB will aspire to review and update this policy accordingly. However, until the ICB adopts a revised policy, this policy will remain in force and any references in it to NICE guidance will remain valid as far as the decisions of this ICB are concerned.

## 10. References

- National Institute for Health and Care Excellence. *NICE CKS Haemorrhoids Management*. 2021 07/2021 [cited 2024 May]; Available from: <https://cks.nice.org.uk/topics/haemorrhoids/management/management/>.
- England, N., *Haemorrhoid Surgery. Evidence-Based Interventions: Guidance for CCGs 2019*: <https://ebi.aomrc.org.uk/resources/>. p. 28-29.
- Watson, A.J., et al., *A pragmatic, multicentre, randomised controlled trial comparing stapled haemorrhoidopexy to traditional excisional surgery for haemorrhoidal disease (eTHoS): study protocol for a randomised controlled trial*. *Trials*, 2014. **15**: p. 439.
- Watson, A.J., et al., *Comparison of stapled haemorrhoidopexy with traditional excisional surgery for haemorrhoidal disease (eTHoS): a pragmatic, multicentre, randomised controlled trial*. *Lancet*, 2016. **388**(10058): p. 2375-2385.
- Brown, S., et al., *The HubBLE Trial: haemorrhoidal artery ligation (HAL) versus rubber band ligation (RBL) for symptomatic second- and third-degree haemorrhoids: a multicentre randomised controlled trial and health-economic evaluation*. *Health Technol Assess*, 2016. **20**(88): p. 1-150.
- Academy of Medical Royal Colleges. *Haemorrhoid surgery*. 2024 [cited 2024; Available from: <https://ebi.aomrc.org.uk/interventions/haemorrhoid-surgery/>.
- Thaha, M.A. and R.J.C. Steele, *BMJ Best Practice Haemorrhoids*. 2022: <https://bestpractice.bmj.com/topics/en-gb/181/pdf/181/Haemorrhoids.pdf>.
- van Tol, R.R., et al., *European Society of ColoProctology: guideline for haemorrhoidal disease*. *Colorectal Dis*, 2020. **22**(6): p. 650-662.
- Hawkins, A.T., et al., *The American Society of Colon and Rectal Surgeons Clinical Practice Guidelines for the Management of Hemorrhoids*. *Dis Colon Rectum*, 2024. **67**(5): p. 614-623.

- El-Kelani, M.Z., et al., *Recommendations and best practice on the management of hemorrhoidal disease in Saudi Arabia*. Hospital Practice, 2022. **50**(2): p. 104-109.
- De Schepper, H., et al., *Belgian consensus guideline on the management of hemorrhoidal disease*. Acta Gastroenterol Belg, 2021. **84**(1): p. 101-120.
- Gallo, G., et al., *Consensus statement of the Italian society of colorectal surgery (SICCR): management and treatment of hemorrhoidal disease*. Techniques in Coloproctology, 2020. **24**(2): p. 145-164.
- National Institute for Health and Care Excellence, *IPG627 Superior rectal artery embolisation for haemorrhoids*. 2018: <https://www.nice.org.uk/guidance/ipg627/resources/superior-rectal-artery-embolisation-for-haemorrhoids-pdf-1899873987681733>.
- National Institute for Health and Care Excellence, *IPG589 Radiofrequency treatment for haemorrhoids*. 2017: <https://www.nice.org.uk/guidance/ipg589/resources/radiofrequency-treatment-for-haemorrhoids-pdf-1899872177055685>.
- National Institute for Health and Care Excellence, *IPG525 Electrotherapy for the treatment of haemorrhoids*. 2015: <https://www.nice.org.uk/guidance/ipg525/resources/electrotherapy-for-the-treatment-of-haemorrhoids-pdf-1899871807540165>.
- National Institute for Health and Care Excellence, *MIB201 eXroid for internal haemorrhoids*. 2019: <https://www.nice.org.uk/advice/mib201/resources/exroid-for-internal-haemorrhoids-pdf-2285965386043333>.
- National Institute for Health and Care Excellence, *IPG342 Haemorrhoidal artery ligation*. 2010: <https://www.nice.org.uk/guidance/ipg342/resources/haemorrhoidal-artery-ligation-pdf-1899867569868997>.
- National Institute for Health and Care Excellence, *TA128 Stapled haemorrhoidopexy for the treatment of haemorrhoids*. 2007: <https://www.nice.org.uk/guidance/ta128/resources/stapled-haemorrhoidopexy-for-the-treatment-of-haemorrhoids-pdf-82598140063429>.
- National Institute for Health and Care Excellence, *IPG34 Circular stapled haemorrhoidectomy*. 2003: <https://www.nice.org.uk/guidance/ipg34/resources/circular-stapled-haemorrhoidectomy-pdf-52774099025605>.
- Gallo, G., et al., *Sclerotherapy for hemorrhoidal disease: systematic review and meta-analysis*. Tech Coloproctol, 2024. **28**(1): p. 28.
- Ding, Z., et al., *Rubber band ligation versus coagulation for the treatment of haemorrhoids: a meta-analysis of randomised controlled trials*. Postgraduate Medical Journal, 2023. **99**(1175): p. 946-953.
- Yuan, X.G., et al., *Comparison of the efficacy and safety of different surgical procedures for patients with hemorrhoids: a network meta-analysis*. Tech Coloproctol, 2023. **27**(10): p. 799-811.
- Patel, J., et al., *HEmoRhoidal disease management with Band ligation versus polidocanol Sclerotherapy: a systematic review and meta-analysis (the HerBS Review)*. Int J Colorectal Dis, 2023. **38**(1): p. 112.
- Lohsiriwat, V., et al., *Recurrence Rates and Pharmacological Treatment for Hemorrhoidal Disease: A Systematic Review*. Advances in Therapy, 2023. **40**(1): p. 117-132.
- Tan, V.Z.Z., et al., *Systematic review and meta-analysis of postoperative pain and symptoms control following laser haemorrhoidoplasty versus Milligan-Morgan haemorrhoidectomy for symptomatic haemorrhoids: a new standard*. Int J Colorectal Dis, 2022. **37**(8): p. 1759-1771.
- Grossi, U., et al. *Surgical Management of Hemorrhoidal Disease in Inflammatory Bowel Disease: A Systematic Review with Proportional Meta-Analysis*. Journal of

Clinical Medicine, 2022. **11**, DOI: 10.3390/jcm11030709.

- Trigui, A., et al., *Laser utility in the treatment of hemorrhoidal pathology: a review of literature*. Lasers in Medical Science, 2022. **37**(2): p. 693-699.
- Aibuedefe, B., et al., *An update on surgical treatment of hemorrhoidal disease: a systematic review and meta-analysis*. International Journal of Colorectal Disease, 2021. **36**(9): p. 2041-2049.
- Ruan, Q.Z., et al., *A systematic review of the literature assessing the outcomes of stapled haemorrhoidopexy versus open haemorrhoidectomy*. Tech Coloproctol, 2021. **25**(1): p. 19-33.
- Longchamp, G., et al., *Non-excisional laser therapies for hemorrhoidal disease: a systematic review of the literature*. Lasers in Medical Science, 2021. **36**(3): p. 485-496.
- Sheikh, P., V. Lohsiriwat, and Y. Shelygin, *Micronized Purified Flavonoid Fraction in Hemorrhoid Disease: A Systematic Review and Meta-Analysis*. Advances in Therapy, 2020. **37**(6): p. 2792-2812.
- Zhang, G., et al., *Network meta-analysis of randomized controlled trials comparing the procedure for prolapse and hemorrhoids, Milligan-Morgan hemorrhoidectomy and tissue-selecting therapy stapler in the treatment of grade III and IV internal hemorrhoids(Meta-analysis)*. International Journal of Surgery, 2020. **74**: p. 53-60.
- Du, T., et al., *Comparison of surgical procedures implemented in recent years for patients with grade III and IV hemorrhoids: a network meta-analysis*. International Journal of Colorectal Disease, 2019. **34**(6): p. 1001-1012.

## 11. Associated OPCS/ICD codes

|  |
|--|
| <b>OPCS codes</b>  |
| H511, H512, H513, H518, H519, H521, H522, H523, H524, H528, H529, H531, H532, H533, H538, H539, L703, Y524, Y532, Z378 - Haemorrhoid Surgery |
| <b>ICD-10 codes</b>  |
| K640, K641, K642, K643, K644, K645, K648, K649 - Haemorrhoids  |

## Appendix 1: Definitions and classifications

### Classification of haemorrhoids<sup>1</sup>

**External haemorrhoids:** Haemorrhoids that are located in the distal anal canal, below the dentate line, and covered by sensitive lining (epithelium) of the anus or skin.

**Internal haemorrhoids:** Haemorrhoids that originate above to the dentate line and covered by insensitive transitional epithelium. Table 1 shows the classification of internal haemorrhoids. This grading of internal haemorrhoids is only a reflection of the degree of prolapse but not a measure of either the disease severity or of the size of hemorrhoidal prolapse.

*Table 1: Classification of internal haemorrhoids*

| Grade   | Description   |
|---------|---|
| Grade 1 | Haemorrhoid protrusion is limited to within the anal canal                                      |
| Grade 2 | Haemorrhoid protrudes beyond the anal canal but spontaneously reduces on cessation of straining |
| Grade 3 | Haemorrhoid protrudes outside the anal canal and reduces fully on manual pressure               |
| Grade 4 | Haemorrhoid protrudes outside the anal canal and is irreducible                                 |

**Mixed haemorrhoids:** These are located both above and below the dentate line.<sup>23</sup>

#### **Persistent haemorrhoid:**

These are haemorrhoid whose symptoms failed to resolve spontaneously or improve with dietary changes, banding or perhaps in certain cases injection. <sup>1</sup>

#### **Recurrent haemorrhoid:**

These are haemorrhoids that have redeveloped following successful treatment.